arXiv:1211.1715v1 [math.SG] 7 Nov 2012

EXACT LAGRANGIAN IMMERSIONS WITH ONE DOUBLE POINT REVISITED

TOBIAS EKHOLM AND IVAN SMITH

Abstract. We study exact Lagrangian immersions with one double point of a closed orientable manifold K into $\mathbb{C}^n$. Our main result is that if the Maslov grading of the double point does not equal 1 then K is homotopy equivalent to the sphere, and if, in addition, the Lagrangian Gauss map of the immersion is stably homotopic to that of the Whitney immersion, then K bounds a parallelizable $(n+1)$-manifold. The hypothesis on the Gauss map always holds when $n = 2k$ or when $n = 8k - 1$. The argument studies a filling of K obtained from solutions to perturbed Cauchy-Riemann equations with boundary on the image $f(K)$ of the immersion. This leads to a new and simplified proof of some of the main results of [11], which treated Lagrangian immersions in the case $n = 2k$ by applying similar techniques to a Lagrange surgery of the immersion, as well as to an extension of these results to the odd-dimensional case.

1. Introduction

We consider Euclidean space $\mathbb{C}^n$ with coordinates $(x_1 + iy_1, \ldots, x_n + iy_n)$ and with its standard exact symplectic structure $\omega = -d\theta$, where $\theta = \sum_{j=1}^{n} y_j\, dx_j$. We are concerned with exact Lagrangian immersions $f\colon K \to \mathbb{C}^n$ of closed orientable manifolds K, i.e. immersions for which $f^*\theta = dz$ for a smooth function $z\colon K \to \mathbb{R}$. Gromov [17] proved that $\mathbb{C}^n$ contains no embedded exact Lagrangian submanifolds, so immersions with a single (transverse) double point have the simplest possible self-intersection.

This paper is a sequel to [11], which shows that if $n > 4$ is even, if $\chi(K) \neq -2$, and if K admits an exact Lagrangian immersion into $\mathbb{C}^n$ with only one double point, then K is diffeomorphic to the standard n-sphere. The proof of this theorem relied on constructing a parallelizable manifold $\mathcal{B}$ with $\partial\mathcal{B} = K$ from moduli spaces of Floer holomorphic disks with boundary in a Lagrange surgery of the image $f(K)$ of the immersion. The corresponding Lagrange surgery is non-orientable when n is odd, hence cannot bound any parallelizable manifold. In this paper, we study moduli spaces of holomorphic disks with boundary on $f(K)$ that admit a continuous lift into K. This turns out to be analytically somewhat simpler, and leads to a result valid in all dimensions.

To state the theorem, recall that the map $\tilde{f} = (f, z)\colon K \to \mathbb{C}^n \times \mathbb{R}$ is a Legendrian immersion (generically an embedding) into $\mathbb{C}^n \times \mathbb{R}$ with the contact form $dz - \theta$. The double points of f correspond to Reeb chords of $\tilde{f}$, and as such have a mod 2 Maslov grading $|\cdot|_2$, which lifts to a mod $2j$ grading $|\cdot|_{2j}$ if the Maslov index of f equals $2j$. (Note that the Maslov index is even since K is orientable). Here $|\cdot|_0 = |\cdot|$ is to be understood as an integer grading. The Lagrangian immersion also determines a Gauss map $K \to U_n/O_n$, taking $p \in K$ to the tangent plane $df(T_pK)$ viewed

---

TE was partially supported by the Göran Gustafsson Foundation for Research in Natural Sciences and Medicine. IS was partially supported by European Research Council grant ERC-2007-StG-205349.

as a Lagrangian subspace of $\mathbb{C}^n$. The *stable Gauss map* Gf is the composition of the Gauss map with the natural map $U_n/O_n \to U/O$. Finally, we recall the classical Whitney immersion in Definition 5.2.

Theorem 1.1. *Let K be an orientable n-manifold that admits an exact Lagrangian immersion f into $\mathbb{C}^n$ of Maslov index $2j$, with exactly one double point a. Suppose $|a|_{2j} \neq 1$. Then $j = 0$, $|a| = n$, and $K \simeq S^n$ is homotopy equivalent to the sphere. If in addition the stable Gauss map Gf of K is homotopic to the stable Gauss map of the Whitney immersion of S^n then K bounds a parallelizable manifold.*

The stable Gauss map Gf of a Lagrangian immersion f of a homotopy n-sphere lifts to a map into U, hence vanishes if n is even. If n is odd it does not necessarily vanish; the relevant homotopy groups are

$$\pi_n(U/O) = \begin{cases} 0 & \text{for } n = 8k-1, \\ \mathbb{Z} & \text{for } n = 8k+1 \text{ or } n = 8k+5, \\ \mathbb{Z}_2 & \text{for } n = 8k+3. \end{cases}$$

In particular, the homotopy condition in Theorem 1.1 always holds when $n = 8k - 1$.

The only even-dimensional homotopy sphere which bounds a parallelizable manifold is the standard sphere, by [20]. If n is even, the mod 2 Maslov grading of the double point is determined by whether the Euler characteristic is $+2$ or -2. Combining these facts, Theorem 1.1 in particular recovers the results on even dimensional spheres in [11] as stated above.

Theorem 1.1 should be contrasted with the results of [12], which provide general constructions of exact Lagrangian immersions with few double points, either by appeal to an h-principle for loose Legendrian embeddings and their Lagrangian caps [23, 13] or via higher-dimensional analogues of the explicit constructions of [27]. In particular, special cases of the result of [12] show that if K is an orientable n-manifold with $TK \otimes \mathbb{C}$ trivial, then

- if $n = 2k$ and $\chi(K) = -2$, then K admits an exact Lagrangian immersion with exactly one double point;
- if $n = 2k + 1$ and K is an odd-dimensional homotopy n-sphere, then K admits an exact Lagrangian immersion with exactly one double point.

Thus, the main result of [11] is essentially sharp, and the Maslov index hypothesis is crucial for the rigidity results established in Theorem 1.1.

In the proof of Theorem 1.1 the homotopy type of K is controlled by a version of linearized Legendrian contact homology adapted to an abstract covering space of K, analogous to Damian's lifted Floer homology [4]. For a more systematic development of that subject, see [14]. The essential simplification in the subsequent construction of a bounding manifold for K, as compared to [11], comes from the fact that breaking of Floer holomorphic disks is concentrated at the double point, and therefore Gromov-Floer boundaries of relevant moduli spaces are ordinary rather than fibered products.

If $n = 4k + 1$ there is at most one exotic sphere that bounds a parallelizable manifold. In dimensions $n = 4k - 1$ there are typically many such exotic spheres, distinguished by the possible signatures of parallelizable $4k$-manifolds which they bound. This paper closes with a brief discussion of the signatures of the bounding manifolds obtained by Floer theory, see Section 5.5.

Finally, we point out that [11] also gave applications to Arnold's "nearby Lagrangian submanifold conjecture", proving that $T^*((S^{n-1}\times S^1)\#\Sigma)$ and $T^*(S^{n-1}\times S^1)$ are not symplectomorphic for a non-trivial homotopy n-sphere Σ when $n=8k$. That result relies essentially on the appearance of the Lagrange surgery of $f(K)$, and does not seem accessible via the simplified argument of this paper.

The paper is organized as follows: in Section 2 we derive the homotopy theoretic conclusions of Theorem 1.1. In Section 3 we discuss aspects of moduli spaces of Floer disks and in Section 4 we establish gluing results that give C^1-parametrizations of neighborhoods of the boundary of these disk moduli spaces. Finally, in Section 5 we construct a filling manifold and prove it is parallelizable. To avoid special cases, we assume throughout the paper that the dimension n of the source of the Lagrangian immersion satisfies $n>2$. Many of the arguments in the paper are close cousins of corresponding arguments in [11], and in such places our treatment is accordingly relatively brief.

2. Topological constraints on exact immersions

We begin with some background material on Legendrian contact homology. For details, see [8, 10] and references therein.

Let $\Lambda\subset(\mathbb{C}^n\times\mathbb{R}, dz-\theta)$ be an orientable Legendrian submanifold. There is an associated Floer-theoretic invariant $LH(\Lambda;k)$, the *Legendrian contact homology* of Λ with coefficients in the field k. This is a unital differential graded algebra generated by the Reeb chords of Λ, and with differential counting certain punctured holomorphic curves in $\mathbb{C}^n$ with boundary on the immersed Lagrangian projection of Λ. In the situations studied in this paper, all Reeb chords have non-zero length, and hence the algebra generators can be canonically identified with the double points of the Lagrangian projection. An *augmentation* of $LH(\Lambda;k)$ is a k-linear dga-map $\epsilon\colon LH(\Lambda;k)\to k$, where the field is concentrated in degree 0 and equipped with the trivial differential. Such a map leads to a linearization of the the dga as follows. The k-vector space underlying the chain complex $C=C^{\mathrm{lin}}(\Lambda;k)$ of the linearization is generated by the Reeb chords of Λ. To define the differential on C, consider the tensor algebra

$$\overline{C} \;=\; k\;\oplus\; C\;\oplus\; C\otimes C\;\oplus\;\cdots\;\oplus\; C^{\otimes m}\;\oplus\;\cdots$$

and view the dga differential ∂ as an element

$$\partial\in C^*\otimes\overline{C},\quad \partial=\sum_{\text{Reeb chords } a} a^*\otimes(\partial a),$$

where C^* is the dual of C and where we view (∂a) as a linear combination of monomials. There is a canonical identification between C and its space of first-order variations. With this in mind, the differential d on C is the element in $C^*\otimes C$ given by composing the first order variation $\delta\partial$ with the augmentation map on coefficients. More concretely, if

$$\partial=\sum_{i;j_1,\dots,j_n}\kappa_{i;j_1,\dots,j_n}\, a_i^*\otimes b_{j_1}\dots b_{j_n},\quad \kappa_{i;j_1,\dots,j_n}\in k$$

then

$$\delta\partial=\sum_{i;j_1,\dots,j_n}\kappa_{i;j_1,\dots,j_n}\, a_i^*\otimes\big((\delta b_{j_1})b_{j_2}\dots b_{j_n}+\cdots+b_{j_1}\dots b_{j_{n-1}}(\delta b_{j_n})\big)$$

and

$$d = \sum_{i;j_1,\dots,j_n} \kappa_{i;j_1,\dots,j_n}\, a_i^* \otimes \big(\epsilon(b_{j_2}\dots b_{j_n})\, b_{j_1} + \dots + \epsilon(b_{j_1}\dots b_{j_{n-1}})\, b_{j_n}\big) \in C^* \otimes C.$$

The invariant $LH(\Lambda;k)$ is well-defined as a mod 2 graded algebra for arbitrary Λ if the characteristic of k equals 2; in this generality, linearizations should preserve the mod 2 degree. If Λ admits a spin structure, then one can work with fields k of arbitrary characteristic. If Λ has vanishing Maslov class, then the mod 2 grading can be refined to a $\mathbb{Z}$-grading, and one can ask for graded linearizations.

Suppose $\pi\colon \tilde{\Lambda} \to \Lambda$ is an abstract (not necessarily finite degree) connected covering space of Λ and that $\epsilon\colon LH(\Lambda;k) \to k$ is an augmentation. We will use a variant chain complex $\widetilde{C}^{\mathrm{lin}}(\tilde{\Lambda},\pi;k)$ which takes account of the cover, and which is analogous to the "lifted Floer homology" of Damian [4]. To define it we first pick a base point $x \in \Lambda$ and pick "capping paths" connecting all Reeb chord endpoints to x. Then, as in [10], consider the Legendrian $\Lambda \cup \Lambda'$ comprising two copies of Λ, the original one and one copy Λ' displaced by a large distance in the Reeb direction. This is a Bott-degenerate situation: through each point of Λ there is a Reeb chord ending on Λ'. We perturb by fixing a sufficiently C^2-small Morse function on Λ and replace Λ' by the graph of the 1-jet of this function, in a suitable Weinstein tubular neighborhood $\approx J^1(\Lambda)$ of Λ'. We keep the notation Λ' for this perturbed component.

The chain complex $\tilde{C} = \widetilde{C}^{\mathrm{lin}}(\tilde{\Lambda},\pi;k)$ is generated by the preimages under π of all endpoints of mixed Reeb chords, i.e. chords that start on Λ and end on Λ'. Let $\tilde{c}$ denote such a preimage. The differential $\tilde{d}\tilde{c}$ is defined as follows. Let c be the mixed Reeb chord corresponding to $\tilde{c}$ and consider a rigid holomorphic disk with boundary on $\Lambda \cup \Lambda'$, with positive puncture at c, and with exactly one mixed negative puncture. Let the augmentation ϵ act on all pure negative punctures. We say that such a disk is *contributing* if ϵ takes non-zero values at all pure Reeb chords along its boundary, otherwise we say it is *non-contributing.* For contributing disks we add the capping paths at all pure Reeb chord endpoints in Λ. In this way, any contributing disk gives a path in Λ; given the start point $\tilde{c} \in \tilde{\Lambda}$ that path determines a unique path in $\tilde{\Lambda}$, which ends at a point $\tilde{b}$ which is one of the preimages of the endpoint of the mixed negative puncture in the disk. We define the *coefficient* of any contributing disk to be the product of the values of ϵ on its pure chords. Finally, the differential $\tilde{d}\tilde{c}$ is defined as the sum of $\tilde{b}$ over all such rigid disks, weighted by the corresponding coefficient.

Remark 2.1. We point out that even when the covering degree and chain groups are infinite-rank, the differential of any given generator is well-defined (i.e. involves only finitely many terms), since the set of chords of action bounded below by some fixed value is always finite, compare to [2, Lemma 2.1]. Note that the chain complex $\tilde{C}$ depends on the choice of capping paths.

For the purposes of this paper, we will need only a number of elementary properties of this theory which we discuss next. We refer to [14] for a more systematic study of the construction. Note first that the mixed Reeb chords of $\Lambda \cup \Lambda'$ are of three kinds: short, Morse, and long. Here short chords correspond to chords that start near the upper endpoint of a Reeb chord of Λ and end near the lower end point of a Reeb chord of Λ', Morse chords correspond to critical points of the small perturbing function, and long chords start near the lower end point of a Reeb chord in Λ and end near the upper endpoint of a Reeb chord of Λ'. Let $\tilde{S}$, $\tilde{C}_{\mathrm{Morse}}$, and $\tilde{L}$ denote the subspaces of $\tilde{C}$ generated by preimages of short, Morse, and long chords, respectively. Since any holomorphic disk has positive area, one finds that

$$\tilde{S} \quad \text{and} \quad \tilde{S} \oplus \tilde{C}_{\text{Morse}}$$

are both sub-complexes of $\tilde{C}$.

Assume now that Λ has vanishing Maslov class. There is then a grading on $\tilde{C}$ that is well-defined up to an overall integer shift. In order to fix the grading we fix the Maslov potential difference between Λ' and Λ to be 1. More prosaically, the grading of a chord is defined as a certain Maslov index of its associated capping path. For mixed chords, such capping paths also include a path going between the components Λ, Λ', and we take this path to have Maslov index 1, see [3, Section 2.1]. With this choice, elements $\tilde{c} \in \tilde{C}_{\text{Morse}}$ are graded by the Morse index of the corresponding critical point, whilst elements $\tilde{s} \in \tilde{S}$ and $\tilde{l} \in \tilde{L}$ are graded by shifts of their gradings downstairs: $|\tilde{s}| = |s| - (n+1)$ and $|\tilde{l}| = |l| + 1$, where s and l are the Reeb chords of Λ corresponding to $\tilde{s}$ and $\tilde{l}$, respectively.

Finally, we will use the fact that the homology of $\tilde{C}$ is invariant under Hamiltonian isotopies of $\mathbb{C}^n$ and their induced actions on Λ; this is straightforward to derive from corresponding results for the two-copy Legendrian contact homology established in [6, Section 2].

We now return to our particular situation. Let $f\colon K \to \mathbb{C}^n$ be an exact Lagrangian immersion of an orientable manifold with one double point, and let $\tilde{f}\colon K \to \mathbb{C}^n \times \mathbb{R}$ be the associated Legendrian embedding. Let a denote the unique double point, which is the canonical generator of $LH(\tilde{f}(K); k)$.

Lemma 2.2. *One of the following holds:*

1. *$LH(\tilde{f}(K); k)$ admits no mod 2 linearization. In this case, $|a|_2 = 1$, and if $LH(\tilde{f}(K); k)$ admits an integer grading then $|a| = 1$.*
2. *$LH(\tilde{f}(K); \mathbb{Z}_2)$ admits a (mod 2 graded) linearization. In this case, the Maslov index of f vanishes, $LH(\tilde{f}(K); k)$ admits an integer grading, and $|a| = n$. Furthermore, K is a homotopy sphere.*

Proof. For the first case, any dga generated by a single element a can only fail to be linearizable if $|a|_{2j} = 1$, for any value j for which the dga admits a mod $2j$ grading.

Suppose then $LH(\tilde{f}(K); \mathbb{Z}_2)$ admits a linearization. Since $\tilde{f}(K)$ is displaceable by Hamiltonian isotopy, the double point bounds of [10, 6] apply and show that

$$\operatorname{rank}_{\mathbb{Z}_2}(H^*(K; \mathbb{Z}_2)) \leq 2 \tag{2.1}$$

which implies that K is a $\mathbb{Z}_2$-homology sphere. Therefore $H^2(K; \mathbb{Z}_2) = 0$, so K is spin and $LH(\tilde{f}(K); k)$ can be defined with coefficients in a characteristic zero field k. The double point bound (2.1) over k then implies that K is a rational homology sphere, hence has vanishing Maslov class, and [6, Theorem 5.5] then implies $|a| = n$. Finally, working with finite field co-efficients, one concludes that K is a $\mathbb{Z}$-homology sphere.

It remains to prove that $\pi_1(K) = 1$. Consider a connected covering space $\pi\colon \tilde{K} \to K$ of K. Then the complex $\tilde{C}^{\text{lin}}(\tilde{K}, \pi; k)$ discussed above, has the form

$$\tilde{S} \oplus \tilde{C}_{\text{Morse}} \oplus \tilde{L},$$

where all elements in $\tilde{S}$ have grading -1 and all elements in $\tilde{L}$ grading $n+1$. Hamiltonian displaceability of $f(K)$ implies that the total homology of the complex vanishes. Since $\tilde{K}$ is connected, this in turn implies $|\tilde{S}| = 1$. That implies the covering π has degree 1, and since the covering was arbitrary, that implies that $\pi_1(K) = 1$, as required. $\square$

We should point out that one can also prove Lemma 2.2 by applying the results of [4] to the Lagrange surgery of $f(K)$, but that argument is somewhat more roundabout and involves more heavy algebraic machinery.

3. Moduli spaces and breaking

3.1. **Displacing Hamiltonian and the Floer equation.** For the rest of the paper, we fix a Lagrangian immersion $f\colon \Sigma \to \mathbb{C}^n$ of a homotopy n-sphere Σ with exactly one transverse double point a. The double point has Maslov grading $|a| = n$. As above we write $\tilde{f}\colon \Sigma \to \mathbb{C}^n \times \mathbb{R}$ for the Legendrian lift of f, and we write $a^{\pm} \in \Sigma$ for the upper and lower endpoints of the Reeb chord corresponding to a.

Following [24], consider a compactly supported time-dependent Hamiltonian function $H\colon \mathbb{C}^n \times [0,1] \to \mathbb{R}$ with the following properties:

- there exists $\epsilon_0 > 0$ such that H_t is constant for $t \in [0,\epsilon_0] \cup [1-\epsilon_0, 1]$;
- the time 1 flow ϕ^1 of the Hamiltonian vector field X_H of H displaces $f(\Sigma)$ from itself: $\phi^1(f(\Sigma)) \cap f(\Sigma) = \varnothing$.

We can suppose that $f(\Sigma)$ is real analytic [11, Lemma 4.1], that the double point of f lies at $0 \in \mathbb{C}^n$, that the branches of $f(\Sigma)$ near the double point are flat and co-incide with $\mathbb{R}^n \cup i\mathbb{R}^n$ in a sufficiently small ball around the origin.

We fix a smooth family of smooth functions

$$\alpha_r\colon \mathbb{R} \longrightarrow [0,1] \quad \text{for } r \in [0,\infty)$$

satisfying the conditions

- $\alpha_r(s) \;= \begin{cases} 0 & |s| > r+1 \\ 1 & |s| \le r \end{cases}$
- $\alpha_r'(s)$ is $\begin{cases} < 0 & s \in (r, r+1) \\ > 0 & s \in (-r-1, -r) \end{cases}$
- $\alpha_r(s) = r\alpha_1(s)$ when $s \in [0,1]$, so in particular $\alpha_0 \equiv 0$.

If D denotes the unit disk in $\mathbb{C}$ and $\xi \in S^1$, then there is a unique conformal map

$$\zeta\colon \mathbb{R} \times [0,1] \to D \quad \text{with} \quad \zeta(\pm\infty) = \pm 1 \text{ and } \zeta(0) = -i, \tag{3.1}$$

with inverse

$$\zeta^{-1} = s + it\colon D \to \mathbb{R} \times [0,1]. \tag{3.2}$$

Let $\beta : [0,\infty) \to [0,1]$ be a non-decreasing smooth function which vanishes near 0 and is identically 1 for $r \ge 1$. Write γ_r for the 1-form on D such that

$$\zeta^* \gamma_r = \beta(r)\alpha_r(s)\, dt.$$

Fix a small $\delta > 0$ and let $\mathcal{J}_\Sigma$ denote the space of almost complex structures on $\mathbb{C}^n$ which tame the standard symplectic form and which agree with the standard structure J_0 in a δ-neighborhood of $f(\Sigma)$. Fix a finite set of points $\mathbf{z} = \{z_1, \dots, z_m\} \in \partial D$ subdivided into two subsets $\mathbf{z} = \mathbf{z}_+ \cup \mathbf{z}_-$ and write

$$D_{\mathbf{z}} = D - \mathbf{z}.$$

Furthermore, for $p, q \geq 0$ write $D_{p,q}$ for $D_{\mathbf{z}}$ for an unspecified set $\mathbf{z} = \mathbf{z}_+ \cup \mathbf{z}_-$ where there are p boundary punctures in $\mathbf{z}_+$ and q in $\mathbf{z}_-$. Note that $\partial D_{p,q}$ is a collection of $p + q - 1$ open arcs. For $r \in [0, \infty)$ and $J \in \mathcal{J}_L$ we will consider Floer disks with boundary on $\tilde{f}(\Sigma)$ with p positive and q negative boundary punctures at a. Such a disk consists of the following:

- A map $u\colon (D_{p,q}, \partial D_{p,q}) \to (\mathbb{C}^n, f(\Sigma))$ for which $u|_{\partial D_{p,q}}$ admits a continuous lift $\tilde{u}\colon \partial D_{p,q} \to \tilde{f}(\Sigma)$ with the property that for each positive boundary puncture $\xi \in \mathbf{z}_+$, $\lim_{\zeta\to\xi\pm} \tilde{u}(\zeta) = a^{\mp}$ and for each negative boundary puncture $\eta \in \mathbf{z}_-$, $\lim_{\zeta\to\eta\pm} \tilde{u}(\zeta) = a^{\pm}$. Here $\lim_{\zeta\to z+}$ and $\lim_{\zeta\to z-}$ denote the limits as $\zeta \to z$ in the counterclockwise and clockwise directions, respectively.
- For some $r(u) = r \in [0, \infty)$, the map u solves the perturbed Cauchy-Riemann equation
$$(du + \gamma_r \otimes X_H)^{0,1} = 0, \tag{3.3}$$
where $A^{0,1}$ denotes the (J, i) complex anti-linear part of the linear map $A\colon TD \to T\mathbb{C}^n$.

Remark 3.1. Since H_t is constant near $t = 0, 1$ and since γ_r has compact support, it follows that for each r_0 there exists δ_0 such that $\gamma_r \otimes X_H = 0$ in a δ_0-neighborhood of $\partial D_{p,q}$. Since in addition $J = J_0$ near L, (3.3) reduces to the ordinary $\bar{\partial}$-equation $\bar{\partial}_{J_0} u = du^{0,1} = 0$ in this δ_0-neighborhood. In particular, this holds in a neighborhood of each puncture and we find that each Floer disk admits a Fourier expansion

$$u(\tau + i\sigma) = \sum_{k\geq 0} c_k e^{-(k+\frac{1}{2})\pi(\tau+i\sigma)}, \quad c_k \in \begin{cases} \mathbb{R}^n \text{ for all } k \text{ at a positive puncture} \\ i\mathbb{R}^n \text{ for all } k \text{ at a negative puncture,} \end{cases}$$

where $\tau + it \in [0, \infty) \times [0, 1]$ conformally parameterizes a neighborhood of one of the punctures of $D_{p,q}$.

Let $\ell(a)$ denote the length of the Reeb chord corresponding to the double point a.

Lemma 3.2. *There is a constant $C > 0$ such that for every Floer disk u with p positive and q negative punctures, the following holds:*

$$\|du\|^2_{L^2} \leq C \left((p - q)\ell(a) + \int_0^1 (\max H_t - \min H_t)\, dt \right).$$

Proof. This follows from a well-known application of Stokes theorem. (The value of the constant $C > 0$ is related to the taming condition for J and ω.) □

If $r = 0$, the Hamiltonian term in (3.3) vanishes identically, and Floer holomorphic disks are exactly holomorphic disks, i.e. maps satisfying the unperturbed Cauchy-Riemann equations.

Corollary 3.3. *Suppose $r = 0$. The space of solutions of* (3.3) *with no boundary punctures is canonically diffeomorphic to Σ, viewed as a space of constant maps. Every non-constant solution to* (3.3) *has at least one positive puncture.*

Proof. For the first statement, Lemma 3.2 implies that the derivative of any solution is zero, hence the solutions are all constant maps. The fact that the moduli space of constant maps is transversely cut out and diffeomorphic to Σ is standard, and follows from the surjectivity of the ordinary $\bar{\partial}$-operator for maps $D \to \mathbb{C}^n$ with constant boundary condition $\mathbb{R}^n$. The second part is immediate from Lemma 3.2. □

3.2. **Linearization and transversality.** The linearization of (3.3) is a C^1-Fredholm operator which depends smoothly on the parameters r and $\mathbf{z}$. To make this claim precise, we begin by defining appropriate configuration spaces of maps, following [10]. We briefly recall the set-up.

Fix a family of smooth reference maps $w_{\mathbf{z}} \colon (D_{\mathbf{z}}, \partial D_{\mathbf{z}}) \to (\mathbb{C}^n, f(\Sigma))$ with positive and negative punctures as prescribed; the map $w_{\mathbf{z}}$ should be constantly equal to the double point a in a small neighborhood of each puncture. Consider a metric on $D_{\mathbf{z}}$ which agrees with the standard metric in half-strip coordinates near each puncture. The configuration space $\mathcal{X}_{p,q}$ will then fiber over the space $Z_{p,q} \subset (\partial D)^{p+q}$ of possible configurations of p positive and q negative boundary punctures.

Let $\mathcal{H}(D_{\mathbf{z}}, \mathbb{C}^n)$ denote the Sobolev space of maps having two derivatives in L^2. Let $\mathcal{H}(D_{\mathbf{z}}, \mathbb{C}^n; w_{\mathbf{z}})$ denote the affine Sobolev space of maps $u : D_{\mathbf{z}} \to \mathbb{C}^n$ of the form $u = u' + w_{\mathbf{z}}$, with $u' \in \mathcal{H}(D_{\mathbf{z}}, \mathbb{C}^n)$. The fiber over $\mathbf{z} \in Z_{p,q}$ is the closed Banach submanifold

$$\mathcal{X}_{p,q}(\mathbf{z}) \subset \mathcal{H}(D_{\mathbf{z}}, \mathbb{C}^n; w_{\mathbf{z}})$$

of those maps u that satisfy the boundary conditions, i.e. $u(\partial D_{\mathbf{z}}) \subset f(\Sigma)$ and $u|_{\partial D_{\mathbf{z}}}$ admits a continuous lift, which have the correct behavior near the punctures, and for which the trace the trace $(du)^{0,1}|_{\partial D_{\mathbf{z}}} = 0$. Equation (3.3) can then be viewed as giving a Fredholm section of the bundle $\mathcal{Y}_{p,q} \to \mathcal{X}_{p,q}$ over this configuration space which has fiber at $u \colon D_{p,q} \to \mathbb{C}^n$ equal to the space of (i, J)-complex anti-linear maps $TD_{p,q} \to u^* T\mathbb{C}^n$.

We write $\mathcal{F}^r_{p,q}$ for the space of Floer disks with p positive and q negative punctures satisfying (3.3) with parameter r. Let $\mathcal{F}_{p,q} = \bigcup_{r \in [0,\infty)} \mathcal{F}^r_{p,q}$. Finally, let $\mathcal{M}_{p,q}$ be the moduli space of holomorphic disks (solutions to the unperturbed Cauchy-Riemann equation) with p positive and q negative punctures, modulo automorphism.

Lemma 3.4. *The formal dimension of the solution spaces are the following:*

$$\dim(\mathcal{F}^r_{p,q}) = 2p + (1-q)n, \tag{3.4}$$

$$\dim(\mathcal{F}_{p,q}) = 2p + (1-q)n + 1, \tag{3.5}$$

$$\dim(\mathcal{M}_{p,q}) = (n-3) + 2p - nq. \tag{3.6}$$

Proof. Since Σ is a homotopy sphere the Maslov index of f vanishes. The index of the linearized problem corresponding to a holomorphic disk with a positive puncture at a of grading $|a| = n$ was computed in [8] to be $n+1$. The formulae above then follow from repeated application of the additivity of the index under gluing and the fact that the dimension of the space of conformal structures on a disk with $p+q$ punctures equals $p+q-3$. □

The perturbation theory for obtaining transversality for the spaces $\mathcal{F}_{p,q}$ is straightforward, and gives the following result.

Lemma 3.5. *For generic $J \in \mathcal{J}_\Sigma$ and generic displacing Hamiltonian, $\mathcal{F}_{p,q}$ is a smooth transversely cut out manifold of dimension $2p + (1-q)n + 1$.*

Below we will only use Floer disks with negative punctures; to simplify notation we write $\mathcal{F}_q = \mathcal{F}_{0,q}$. Lemma 3.5 shows that for generic data $\mathcal{F}_q = \varnothing$ for $q > 1$, $\mathcal{F}_1$ is 1-dimensional, and $\mathcal{F}_0$ is $(n+1)$-dimensional.

Corollary 3.3 implies that the space $\mathcal{F}_0^0$ of constant maps is transversely cut out and is the boundary of $\mathcal{F}_0$.

3.3. C^1-structure on $\mathcal{M}_{1,0}$.

Let $\mathcal{M} = \mathcal{M}_{1,0}$ denote the space of holomorphic disks with exactly one positive puncture. Let $G_1 \leq PSL_2(\mathbb{R})$ denote the subgroup of maps fixing $1 \in \partial D$. The space $\mathcal{M}$ is a quotient of a space of parametrized holomorphic disks by the reparametrization action of G_1, hence is not directly a subspace of a Banach space. The existence of a smooth structure on $\mathcal{M}$ follows from a gauge-fixing procedure which we now explain.

Recall from Remark 3.1 that any (Floer) holomorphic disk with a positive puncture at the double point a has a Fourier expansion

$$u(\tau + i\sigma) = \sum_{k \geq 0} c_k e^{-(k+\frac{1}{2})\pi(\tau + i\sigma)}, \quad c_k \in \mathbb{R}^n \tag{3.7}$$

in half-strip coordinates around the puncture (which we take to lie at $1 \in \partial D$). Our next result shows that generically the first Fourier coefficient is non-zero; we remark that the vanishing or non-vanishing of this coefficient is independent of the choice of half-strip neighborhood.

Lemma 3.6. *For generic $J \in \mathcal{J}_\Sigma$ and generic Hamiltonian the following hold.*

- *If $u \in \mathcal{M}$ then the leading Fourier coefficient $c_0 \in \mathbb{R}^n$ of its Fourier expansion near its positive puncture is non-zero.*
- *If $u \in \mathcal{F}_1$ then the leading Fourier coefficient $c_0 \in i\mathbb{R}^n$ of its Fourier expansion near its negative puncture is non-zero.*

Proof. To see the first statement, note that solutions with $c_0 = 0$ lie in a Sobolev space with positive weight slightly larger than $\frac{\pi}{2}$. The Fredholm index of the linearized $\bar{\partial}$-operator on such a weighted Sobolev space equals 1, and hence the formal dimension of the solution space is $1 - 2 = -1$, and the space is generically empty, cf. Lemma 3.5. The second statement is proved in a similar way: solutions with vanishing c_0 belong to a weighted Sobolev space on which the corresponding linearized operator has formal dimension $1 - n < 0$. □

Lemma 3.7. *For generic $J \in \mathcal{J}_\Sigma$ and displacing Hamiltonian, the moduli space $\mathcal{M}$ is a compact C^1-smooth manifold of dimension $n-1$.*

Proof. Lemma 3.6 implies that for $\epsilon > 0$ sufficiently small, the following holds for a map $u\colon D_1 \to \mathbb{C}^n$ representing an element in $\mathcal{M}$. If $p^\pm$ is the point in the boundary lift $\tilde{u}$ of u closest to the puncture which lies in the ϵ-sphere $S_\epsilon(a^\pm) \subset \Sigma$ around $a^\pm$, then $p^\pm$ is a transverse intersection between the boundary ∂D_1 and that sphere. We fix gauge (i.e. associate maps to elements of the quotient space $\mathcal{M}$) by letting the puncture correspond to $1 \in \partial D$ and $p^\pm$ to $\pm i \in \partial D$.

There is a Sobolev space $\mathcal{X}$ of maps $\hat{u}\colon D_{1,\pm i} \to \mathbb{C}^n$ with puncture at 1 and marked points at $\pm i$ which are mapped to $S_\epsilon(a^\pm)$. The space $\mathcal{M}$ is naturally identified with the inverse image of 0 under the C^1-section of $\mathcal{Y} \to \mathcal{X}$, where $\mathcal{Y}$ is the space of complex anti-linear maps, corresponding to the $\bar{\partial}$-operator. Perturbing J close to the positive puncture, see [10, Lemma 4.5 (1)], ensures that this zero-section is transversely cut out. This endows $\mathcal{M}$ with the structure of a C^1-manifold, whose dimension is then given by Lemma 3.4.

By Stokes' theorem, disks in $\mathcal{M}$ have minimal area amongst non-constant holomorphic disks, whence Gromov compactness [25] implies that $\mathcal{M}$ is compact. □

Whilst the closest intersection point between ∂D_1 and $S_\epsilon(a^\pm)$ is transverse, it is not necessarily the unique intersection. There is a slightly stronger statement for maps in $\mathcal{F}_1$: we conclude from Lemma 3.6 that for $\epsilon > 0$ sufficiently small and $u \in \mathcal{F}_1$, the transverse intersection point between $\tilde{u}$ and $S_\epsilon(a^\pm)$ guaranteed by the lemma is in fact the only intersection point. Indeed, if this were not true, then by taking limits one would obtain a solution to a point-constrained problem. However, a point condition at $a^\pm$ decreases formal dimension by $n-1$, so the corresponding space would be empty by Lemma 3.5.

3.4. **Gromov-Floer compactness and consequences.** The results obtained so far are essentially summarized by the following.

Lemma 3.8. *If $J \in \mathcal{J}_\Sigma$ and $H\colon \mathbb{C}^n \times [0,1] \to \mathbb{R}$ are generic, then the space $\mathcal{F}_1$ is a compact C^1-manifold of dimension 1, and $\mathcal{F}_0$ is a non-compact C^1-manifold of dimension $n+1$ with Gromov-Floer boundary diffeomorphic to the product $\mathcal{F}_1 \times \mathcal{M}$.*

Proof. By Gromov compactness, any sequence in $\mathcal{F}_q$ has a subsequence converging to a broken disk in $\mathcal{F}_{q+j}$ with j holomorphic disks from $\mathcal{M}$ attached at negative punctures. By Lemma 3.5, $\mathcal{F}_q = \varnothing$ if $q > 1$. The first statement follows. For the second statement, we first note that the argument just given shows that elements in $\mathcal{F}_1 \times \mathcal{M}$ are the only possible broken configurations in the Gromov Floer boundary. A standard gluing argument, see Section 4, shows that all elements in the product appear as limits of sequences in $\mathcal{F}_0$. □

3.5. **Gauge fixing and Gromov-Floer convergence.** It will be important to have more detailed information on Gromov-Floer convergence of sequences in $\mathcal{F}_0$. Fix small spheres $S_\epsilon(a^\pm)$ as in Section 3.3. For any $M > 0$, the open set

$$U_M = \{u \in \mathcal{F}_0 \colon \sup_{z \in D} |du(z)| > M\}$$

is a neighborhood of the Gromov-Floer boundary of $\mathcal{F}_0$. For $M > 0$ and $u \in \mathcal{F}_0$, let $B^{(1)}(u, M) = |du|^{-1}([M, \infty)) \subset D$ and let $\delta(u, M) = \operatorname{diam}(B^{(1)}(u, M))$.

Lemma 3.9. *Let $\epsilon_M = \sup_{u \in U_M} \delta(u, M)$. Then $\epsilon_M \to 0$ as $M \to \infty$.*

Proof. If not, there is a sequence of disks in $\mathcal{F}_0$ with holomorphic disk bubbles forming at (at least) two points. This contradicts Lemma 3.8. □

It follows by Lemma 3.9 that for each $\delta_0 > 0$ there is $M > 0$ such that for every $u \in U_M$, $\delta(u, M) < \delta_0$. Let $I_u \subset \partial D$ be the smallest interval containing $B^{(1)}(u, M) \cap \partial D$ and let ζ_u denote the midpoint of I_u.

Lemma 3.10. *For all sufficiently large $M > 0$, if $u \in U_M$ then there are points $\zeta_\pm$ in I_u such that $u(\zeta_\pm) \in S_\epsilon(a^\pm)$.*

Proof. If the lemma does not hold then there is $\sigma \in \{+, -\}$ and a sequence of solutions $u_j \in U_{M_j}$ with $M_j \to \infty$ such that $u_j(I_{u_j}) \cap S_\epsilon(a^\sigma) = \varnothing$. Passing to a subsequence, we may assume that $\zeta_{u_j} \to \zeta$ converge. By Gromov-Floer compactness, a subsequence of u_j converges to a solution $\tilde{u}$ in $\mathcal{F}_1$ on compacts subsets of $D - \zeta$. It follows that the intersection number of $\tilde{u}(\partial D - \zeta)$ and $S_\epsilon(a^\sigma)$ equals ± 1. Since the intersection number of any closed curve in Σ and $S_\epsilon(a^\sigma)$ equals 0, we find that there is $\zeta_\sigma \in I^{u_j}$ such that $u_j(\zeta_\sigma) \in S_\epsilon(a^\sigma)$ for j large enough. The lemma follows. □

Let H denote the upper half-plane. Given $u \in U_M$, let $H_\delta \subset H$ be a half-disk parameterizing a small neighborhood in D centered on ζ_u. Let $\zeta_\pm$ be the points in ∂H_δ most distant from the origin satisfying $u(\zeta_\pm) \in S_\epsilon(a^\pm)$. Let $\alpha = \frac{\zeta_+ - \zeta_-}{2}$ and $\beta = \frac{\zeta_+ + \zeta_-}{2}$. The map

$$\psi \colon H \to H, \quad \psi(z) = \alpha z + \beta \quad \text{satisfies } \psi(-1) = \zeta_+, \quad \psi(1) = \zeta_-.$$

Lemma 3.9 implies that $\alpha, \beta \to 0$ as $M \to \infty$. Let $\Omega_\rho = \psi^{-1}(H_\delta) \subset H$, a subset of diameter $\rho = \mathcal{O}(1/\delta)$.

Lemma 3.11. *For any sequence of maps $u_l \in U_{M_l}$ with $M_l \to \infty$, the sequence $u_l \circ \psi \colon \Omega_\rho \to \mathbb{C}^n$ has a subsequence that converges uniformly on compact subsets to a holomorphic disk $u \colon (H, \partial H) \to (\mathbb{C}^n, \Sigma)$ with $u(-1) \in S_\epsilon(a^+)$ and $u(1) \in S_\epsilon(a^-)$, which represents an element in $\mathcal{M}$.*

Proof. If the derivative of $u_j \circ \psi$ is uniformly bounded we can extract a convergent subsequence, with limit which must be non-constant since $S_\epsilon(a^+) \cap S(\epsilon)(a^-) = \varnothing$. The lemma then follows from Lemma 3.8.

To see that the derivative must indeed be uniformly bounded, assume not for contradiction. There is at least one point $\xi \in \partial H$ at which a bubble forms, and a rescaling argument as above yields a non-constant holomorphic disk. Lemma 3.8 implies that this is the only bubble and hence $u_j \circ \psi$ must converge to a constant map on compact subsets of $H - \xi$. This however contradicts $S_\epsilon(a^+) \cap S_\epsilon(a^-) = \varnothing$. □

For $\zeta \in \partial D$, let $B(\zeta; \delta)$ denotes a δ-neighborhood of ζ in D.

Corollary 3.12. *For each $\delta > 0$ and $\epsilon_0 > 0$, there exists $M > 0$ and $\rho > 0$ such that for every $u \in U_M \subset \mathcal{F}_0$ there exists $(u_1, u_2) \in \mathcal{F}_1 \times \mathcal{M}$ such that the following holds:*

1. *Let $\zeta \in \partial D$ be the puncture in domain of u_1. The C^1-distance between $u|_{D - B(\zeta;\delta)}$ and $u_1|_{D - B(\zeta;\delta)}$ is smaller than ϵ_0.*
2. *Let $\zeta^\pm$ denote the intersection points of $u_2(\partial H)$ and $S_\epsilon(a^\pm)$ closest to the puncture. Consider the unique representative $u_2 \colon H \to \mathbb{C}^n$ with $u_2(\mp i) = \zeta^\pm \in S_\epsilon(a^\pm)$. The map $u \circ \psi$ is defined on H_ρ and the C^1-distance between $u \circ \psi$ and $u_2|_{H_\rho}$ is smaller than ϵ_0.*

Proof. If (1) or (2) do not hold, then we extract a sequence contradicting Lemma 3.11. □

4. Gluing

The main objective of this section is to establish the following result.

Lemma 4.1. *For generic $J \in \mathcal{J}_L$ and Hamiltonian $H\colon \mathbb{C}^n \times [0,1] \to \mathbb{R}$, the compactified moduli space of Floer disks*

$$\overline{\mathcal{F}}_0 = \mathcal{F}_0 \cup (\mathcal{F}_1 \times \mathcal{M})$$

admits the structure of a C^1-manifold with boundary, $\partial\overline{\mathcal{F}}_0 = \Sigma \sqcup (\mathcal{F}_1 \times \mathcal{M})$.

To prove Lemma 4.1, following Floer's classical gluing argument, we will define a C^1-map parameterizing a neighborhood of the Gromov-Floer boundary of $\mathcal{F}_0$. The construction here is somewhat simpler than its counterpart in [11].

Our main technical tool is [11, Lemma 6.1], which is a version of Floer's Picard lemma [16] that we restate here for convenience. Let T and M be finite dimensional smooth manifolds and let $\pi_X\colon X \to M \times T$ be a smooth bundle of Banach spaces over $M \times T$. Let $\pi_Y\colon Y \to T$ be a smooth bundle of Banach spaces over T. Let $f\colon X \to Y$ be a smooth bundle map of bundles over T and write $f_t\colon X_t \to Y_t$ for the restriction of f to the fiber over $t \in T$ (where the fiber X_t is the bundle over M with fiber $X_{(m,t)}$ at $m \in M$). If $m \in M$, then write $d_m f_t\colon X_{(m,t)} \to Y_t$ for the differential of f_t restricted to the vertical tangent space of X_t at $0 \in X_{(m,t)}$, where this vertical tangent space is identified with the fiber $X_{(m,t)}$ itself; similarly, the tangent spaces of Y_t are identified with Y_t using linear translations. Denote by 0_Y the 0-section in Y, and by $D(0_X;\epsilon)$ an ϵ-disk sub-bundle of X.

Lemma 4.2. *Let $f\colon X \to Y$ be a smooth Fredholm bundle map of T-bundles, with Taylor expansion in the fiber direction:*

$$f_t(x) = f_t(0) + d_m f_t\, x + N_{(m,t)}(x), \quad \text{where } \ 0, x \in X_{(m,t)}. \tag{4.1}$$

Assume that $d_m f_t$ is surjective for all $(m,t) \in M \times T$ and has a smooth family of uniformly bounded right inverses $Q_{(m,t)}\colon Y_t \to X_{(m,t)}$, and that the non-linear term $N_{(m,t)}$ satisfies a quadratic estimate of the form

$$\|N_{(m,t)}(x) - N_{(m,t)}(y)\|_{Y_t} \le C\|x-y\|_{X_{(m,t)}}(\|x\|_{X_{(m,t)}} + \|y\|_{X_{(m,t)}}), \tag{4.2}$$

for some constant $C > 0$. Let $\ker(df_t) \to M$ be the vector bundle with fiber over $m \in M$ equal to $\ker(d_m f_t)$. If $\|Q_{(m,t)} f_t(0)\|_{X_{(m,t)}} \le \frac{1}{8C}$, then for $\epsilon < \frac{1}{4C}$, $f^{-1}(0_Y) \cap D(0_X;\epsilon)$ is a smooth submanifold diffeomorphic to the bundle over T with fiber at $t \in T$ the ϵ-disk bundle in $\ker(df_t)$.

Proof. This is proved using Newton iteration, see [11, Lemma 6.1], where it is also shown that there is a constant κ such that the distance between the initial point of the iteration and the actual solution is controlled by $\kappa\|f_t(0)\|$. □

4.1. **Pre-gluing maps of disks.**

Consider $(u_1,u_2) \in \mathcal{F}_1 \times \mathcal{M}$, where $u_1\colon D_\zeta \to \mathbb{C}^n$ for some boundary puncture $\zeta \in \partial D$, and $u_2\colon D_1 \to \mathbb{C}^n$ satisfies the gauge fixing condition $\tilde{u}_2(\pm i) \in S_\epsilon(a^{\mp})$. As above we have half-strip neighborhoods of the punctures in the two disks where the metric of the disk agrees with the standard metric. For simpler formulae we think of the half strip neighborhood of ζ in the domain of u_1 as parameterized by $(-\infty, 0] \times [0,1]$ and that of 1 in the domain of u_2 as parameterized by $[0,\infty) \times [0,1]$. Write H_j for the domain of u_j. For $\rho \geq 2$, let

$$\begin{aligned} H_{1;\rho} &= H_1 - ((-\infty, -\rho) \times [0,1]), \\ H_{2;\rho} &= H_2 - ((\rho, \infty) \times [0,1]), \end{aligned}$$

and define

$$\Delta_\rho = H_1 \#_\rho H_2 = (H_{1;\rho} \cup H_{2;\rho})/\sim, \tag{4.3}$$

where

$$\rho + it \sim -\rho + it, \qquad -\rho + it \in H_1 \text{ and } \rho + it \in H_2.$$

Then D comes equipped with a metric that agrees with the given metrics on $H_{1;\rho}$ and $H_{2,\rho}$ (which are the standard Euclidean metrics on the strip neighborhoods of ∞). Furthermore, $H_1 \#_\rho H_2$ contains the finite strip region in the middle,

$$[-\rho, \rho] \times [0,1] \approx [0,\rho] \times [0,1] \cup [-\rho, 0] \times [0,1] \subset H_1 \#_\rho H_2. \tag{4.4}$$

Note that for $\rho > 0$ sufficiently large, u_j maps $H_j - H_{j;\rho}$ into a fixed small neighborhood of the double point a of $f(\Sigma)$, where the Lagrangian agrees with $\mathbb{R}^n \cup i\mathbb{R}^n$. For such ρ, there is a preglued map

$$w_\rho = u_1 \#_\rho u_2 \colon (H_1 \#_\rho H_2, \partial(H_1 \#_\rho H_2)) \to (\mathbb{C}^n, f(\Sigma)),$$

which agrees with u_j on $H_{j;\rho-1}$ and linearly interpolates between the two maps in the remaining square inside the standard neighborhood of a. Note that this interpolated map, when restricted to the boundary, has a natural lift into $\tilde{f}(\Sigma)$.

The space of disks D_ζ with one boundary puncture is canonically identified with the boundary circle of the unit disk ∂D. The space of preglued domains Δ_ρ obtained by joining a disk with boundary puncture at 1 to a disk with boundary puncture at an arbitrary point at distance $\rho \geq \rho_0$ in a fixed cylindrical end is then canonically identified with $S^1 \times [\rho_0, \infty)$. There are two marked points $\zeta^\pm$ on the boundary of Δ_ρ corresponding to $\pm i \in D_1 \approx H_2$.

Let $\widehat{\mathcal{X}}$ denote the bundle over $S^1 \times [\rho_0, \infty)$ with fiber over (ρ, q) the space of maps $u \colon \Delta_\rho \to \mathbb{C}^n$ with two derivatives in L^2, which are holomorphic on the boundary, and which admit a continuous boundary lift into Σ taking $\zeta^\pm$ into $S_\epsilon(a^\mp)$.

Lemma 4.3. *$\widehat{\mathcal{X}}$ is a locally trivial C^1-smooth Banach bundle.*

Proof. Local trivializations of $\widehat{\mathcal{X}}$ over $S^1 \times [\rho_0, \infty)$ are given as follows, see [11, Section 4.4] for a more detailed description.

- In the S^1-direction, we precompose with a rotation which moves the puncture in the domain of the first strip H_1, and which does not change the map on $H_{2;\rho}$ but glues this in at the new location;
- in the $[\rho_0, \infty)$-direction we precompose with diffeomorphisms $\psi \colon \Delta_\rho \to \Delta_{\rho'}$ which are the identity outside the region (4.4), and inside that region are perturbations of diffeomorphisms that stretch or shrink the square $[-\rho, \rho] \times [0,1]$ in the first coordinate (but which have been deformed to be holomorphic along the boundary).

Let $\widehat{\mathcal{X}}_\rho$ denote the fiber of $\widehat{\mathcal{X}}$ over $\rho \in [\rho_0, \infty)$, and fix a map $u \in \widehat{\mathcal{X}}_\rho$. Let $\mathcal{E}(u)$ denote the space of vector fields v along u with two derivatives in L^2 that are holomorphic on the boundary, $(\nabla v)^{0,1}|_{\partial\Delta_\rho} = 0$, and that vanish in the direction perpendicular to $S_\epsilon(a^\mp)$ at $\zeta^\pm$. Let $V(u)$ be a 1-dimensional space spanned by a vector field as above that has normal component at $\zeta^\pm$ corresponding to a motion of both marked points in the positive direction, corresponding to the linearization of the ∂D direction.

The tangent space of the fiber $\widehat{\mathcal{X}}_\rho$ at u splits as $\mathcal{E}(u) \oplus V(u)$. Furthermore, there is an exponential map $\exp_u \colon \mathcal{E}(u) \oplus V(u) \to \widehat{\mathcal{X}}_\rho$ which can be written

$$\exp_u(v, c) = \operatorname{Exp}_u(v) \circ r_c^{-1},$$

where $r_c \colon \Delta_\rho \to \Delta'_\rho$ is the map which is a rotation on $H_{1;\rho} \subset D_\zeta$, taking it to $H'_{1;\rho} \subset D_{e^{ic}\zeta}$, and maps $H_{2;\rho}$ canonically to $H'_{2;\rho}$. This gives a map

$$\Delta_\rho = H_{1;\rho} \# H_{2;\rho} \to H'_{1;\rho} \# H'_{2;\rho} = \Delta'_\rho.$$

which induces a C^1-chart on $\widehat{\mathcal{X}}_\rho$ near u.

It remains to show that such charts vary in a C^1-smooth way with $\rho \in [\rho_0, \infty)$. Consider precomposition with a diffeomorphism $\psi \colon \Delta_\rho \to \Delta_{\rho'}$ that is holomorphic along the boundary. The exponential map above is equivariant under precomposition with ψ:

$$\exp_{u\circ\psi}(v \circ \psi, c) = (\exp_u(v, c)) \circ \psi.$$

Fixing the domain for local trivialization, the linearized variation in the ρ-direction is spanned by a vector field that corresponds to moving the marked points toward each other. □

We will use $\widehat{\mathcal{X}}$ as the source space for the Floer operator. We denote the naturally corresponding target space $\widehat{\mathcal{Y}}$. This space is a locally trivial bundle over $[\rho_0, \infty)$ with fiber at ρ equal to the Sobolev space of complex anti-linear maps $A \in \operatorname{Hom}^{0,1}(T\Delta_\rho, \mathbb{C}^n)$ with one derivative in L^2 and $A|_{\partial\Delta_\rho} = 0$, and with local trivializations in the S^1 and $[\rho_0, \infty)$-directions given by precomposition with the differentials of the diffeomorphisms $\Delta_\rho \to \Delta_{\rho'}$ described above.

4.2. **Pre-gluing as a map and the Floer operator.** Let $\mathcal{N} = \mathcal{F}_1 \times \mathcal{M} \times [\rho_0, \infty)$, and let $\mathcal{N}_\rho = \mathcal{F}_1 \times \mathcal{M}$ thought of as the fiber over ρ of the projection $\mathcal{N} \to [\rho_0, \infty)$. Consider the product $\widehat{\mathcal{X}} \times [0, \infty)$ and define the map

$$\operatorname{PG} \colon \mathcal{N} \to \widehat{\mathcal{X}} \times [0, \infty)$$

as follows:

$$\operatorname{PG}(u_1, u_2, \rho) = (u_1 \#_\rho u_2, r(u_1)),$$

where $r(u_1)$ is the coordinate of the Hamiltonian at u_1, i.e. u_1 solves the Floer equation $(du_1 + \gamma_{r(u_1)} \otimes X_H)^{0,1} = 0$. By compactness of the moduli spaces involved we find that if ρ_0 is sufficiently large then PG is a fiber preserving embedding (as a map of bundles over $[\rho_0, \infty)$). More precisely, the restriction

$$\operatorname{PG}_\rho = \operatorname{PG}|_{\mathcal{N}_\rho} \colon \mathcal{N}_\rho \to \widehat{\mathcal{X}}_\rho \times [0, \infty)$$

is a family of embeddings which depends smoothly on ρ.

Consider the normal bundle $N\mathcal{N}_\rho$ of $\mathcal{N}_\rho = \operatorname{PG}_\rho(\mathcal{N}_\rho) \subset \widehat{\mathcal{X}}_\rho \times [0, \infty)$ as a sub-bundle of the restriction $T_{\mathcal{N}_\rho}\widehat{\mathcal{X}}_\rho \times [0, \infty)$ of the tangent bundle of $\widehat{\mathcal{X}}_\rho \times [0, \infty)$ to $\mathcal{N}_\rho$. The fiber $N_{w_\rho}\mathcal{N}_\rho$ of this normal bundle at $(w_\rho, r) = (u_1\#_\rho u_2, r(u_1)) \in \widehat{\mathcal{X}}_\rho \times [0,\infty)$ is the L^2 complement of the subspace $d\,\mathrm{PG}(T_{(u_1,u_2)}\mathcal{N}_\rho) \subset T_{(w_\rho,r)}\widehat{\mathcal{X}}_\rho \times [0,\infty)$. Here, the L^2 pairing is defined as

$$\langle (v,c)\,,\,(\tilde{v},\tilde{c})\rangle = \langle v\,,\,\tilde{v}\rangle_{\mathcal{E}(w_\rho)} + \langle c\,,\,\tilde{c}\rangle_{\mathbb{R}},$$

where the first summand is the L^2-pairing on the Sobolev space $\mathcal{E}(w_\rho)$ induced by the L^2-pairing on the ambient space and the second is the standard inner product on $\mathbb{R}$ viewed as the tangent space to $[0,\infty)$.

By transversality and a standard linear gluing result, see (5.2) below, $d\,\mathrm{PG}(T_{(u_1,u_2)}\mathcal{N}_\rho) = T_{(w_\rho,r)}(\mathcal{F}_1\times\mathcal{M})_\rho$ is given by

$$T_{(w_\rho,r)}(\mathcal{F}_1\times\mathcal{M})_\rho = \ker(u_1)' \oplus \ker(u_2)'.$$

Here $\ker(u_1)'$ is spanned by vector fields of the form $v'+c'$, where $v+c$ lies in the kernel of the linearized operator at u_i, and where v' is a cut-off version of v and c' is the element in $V(w_\rho)$ with the same constant value as $c \in V(u_1)$. Similarly, $\ker(u_2)'$ is spanned by cut-off versions w' of kernel vector fields w of the linearized operator at u_2.

Since the above constructions are smooth in $\rho \in [\rho_0,\infty)$, we get bundles

$$T_{\mathcal{N}}\widehat{\mathcal{X}} \times [0,\infty) \to [\rho_0,\infty),$$
$$N\mathcal{N} \to [\rho_0,\infty),$$

with local trivializations given by precomposition with the diffeomorphisms $\Delta_\rho \to \Delta_{\rho'}$ discussed above, in the latter case followed by L^2-projection to the normal bundle.

The Floer equation now gives a smooth bundle map $f\colon T_{\mathcal{N}}\widehat{\mathcal{X}}\times[0,\infty) \to \widehat{\mathcal{Y}}$, where $f_\rho\colon T_{\mathcal{N}_\rho}\widehat{\mathcal{X}}_\rho \to \widehat{\mathcal{Y}}_\rho$ is defined as follows. If $w_\rho = u_1\#_\rho u_2 \in \mathcal{N}_\rho$ then for $(v,c) \in T_{w_\rho}\widehat{\mathcal{X}}_\rho \times [0,\infty) = \mathcal{E}(u)\oplus V(u)$:

$$f_\rho(v,c) = \Big(d\exp_{w_\rho}(v,c) + \gamma_{r(u_1)} \otimes X_H(\exp_{w_\rho}(v,c))\Big)^{0,1}. \tag{4.5}$$

Below we will keep the notation f for the restriction $f|_{N\mathcal{N}}$.

4.3. **The gluing map.**

To construct the gluing map

$$\Phi\colon \mathcal{N} \to \mathcal{F}_0$$

which parametrizes a C^1-neighborhood of the Gromov-Floer boundary we will apply Lemma 4.2 to the map $f\colon N\mathcal{N} \to \widehat{\mathcal{Y}}$ defined in (4.5). For $(u_1,u_2,\rho) \in \mathcal{N}$, let $w_\rho = u_1\#_\rho u_2$. In the notation of Lemma 4.2:

- $[\rho_0,\infty)$ corresponds to T, $\mathcal{N}$ corresponds to M;
- $N\mathcal{N}$ corresponds to X, $\widehat{\mathcal{Y}}$ corresponds to Y;
- we use the norm $\|\cdot\|_{N\mathcal{N}_\rho}$ which is the usual Sobolev 2-norm on the vector field component and the standard norm in $\mathbb{R}$ on the linearized rotation component, and the corresponding Sobolev 1-norm for $\|\cdot\|_{\widehat{\mathcal{Y}}_\rho}$;
- the fiber $N_{w_\rho}\mathcal{N}$ of the normal bundle $N\mathcal{N}_\rho$ at w_ρ corresponds to $X_{(m,t)}$;
- w_ρ corresponds to $0 \in X_{(m,t)}$.

Lemma 4.4. *There exists $\gamma > 0$ such that for any $(u_1,u_2,\rho) \in \mathcal{N}$,*

$$\|f_\rho(w_\rho)\|_{\widehat{\mathcal{Y}}_\rho} = \mathcal{O}(e^{-\gamma\rho}).$$

Proof. By Fourier expansion near the punctures, $|u_j^{(k)}(z)| = \mathcal{O}(e^{-(\gamma+\delta)\rho})$, $j = 1, 2$, $k \leq 2$, in the interpolation region. □

Lemma 4.5. *For ρ_0 sufficiently large, the vertical differential*

$$d_{w_\rho} f_\rho \colon N_{w_\rho}\mathcal{N}_\rho \to \widehat{\mathcal{Y}}_\rho$$

is surjective and admits a uniformly bounded right inverse. Moreover, the quadratic estimate (4.2) *for the non-linear term in the Taylor expansion of f_ρ holds.*

Proof. This follows from [10, Proof of Proposition 4.6]. □

Lemmas 4.4 and 4.5 have the following consequence:

Corollary 4.6. *The Newton iteration map with initial values in $\mathcal{N}_\rho = \mathrm{PG}(\mathcal{N}_{\{}\rho\}) \subset N\mathcal{N}_\rho$ gives a C^1-diffeomorphism*

$$\Phi \colon \mathcal{F}_1 \times \mathcal{M} \times [\rho_0, \infty) \to f^{-1}(0),$$

where 0 *denotes the* 0*-section in $\widehat{\mathcal{Y}}$, such that $\Phi(u_1, u_2, \rho)$ limits to a broken curve with components $u_1 \in \mathcal{F}_1$ and $u_2 \in \mathcal{M}$ as $\rho \to \infty$.*

Proof. Immediate from Lemma 4.2. □

4.4. **Parameterizing a neighborhood of the Gromov-Floer boundary.** The final steps of the proof mimic the corresponding steps in [11] very closely. We wish to show that $\exp(f^{-1}(0)) \subset \widehat{\mathcal{X}} \times [0, \infty)$ gives a C^1-collar neighborhood $\approx \mathcal{N}$ of the Gromov-Floer boundary of $\mathcal{F}_0$.

Fix $\rho \in [\rho_0, \infty)$, and let $f^{-1}(0)_\rho = f^{-1}(0) \cap N\mathcal{N}_\rho$. As in [11, Lemma 6.7], we introduce a reparametrization map

$$\Psi_\rho \colon f^{-1}(0)_\rho \to \mathcal{F}_0,$$

which writes the domain D as $H_{1;\rho}$ with the disk $H_{2;\rho}$ attached by a scaling $e^{-2\pi\rho}$. An element $u \in N\mathcal{N}_\rho$ gives a map $\Psi_\rho(u) \colon D \to \mathbb{C}^n$, which is the reparametrization above of the map $\exp(u)$, which was originally defined on Δ_ρ. This map solves the Floer equation if and only if $f(u) = 0$. We let $\Psi \colon f^{-1}(0) \to \mathcal{F}_0$ be the map which equals Ψ_ρ on $f^{-1}(0)_\rho$.

Theorem 4.7. *For ρ_0 sufficiently large, the map $\Psi \colon \mathcal{F}_1 \times \mathcal{M} \times [\rho_0, \infty) \to \mathcal{F}_0$ is a C^1 embedding onto a neighborhood of the Gromov-Floer boundary.*

Proof. This is a direct analogue of [11, Lemma 6.7 & Theorem 6.8]. Briefly, we wish to show that the map Ψ is surjective onto a neighborhood of the Gromov-Floer boundary of $\mathcal{F}_0$. Consider a Gromov-Floer convergent sequence w_j with non-trivial bubbling. Corollary 3.12 implies that

- for any fixed ρ_0, the maps w_j C^1-converge uniformly on compact subsets of the two ends of the strip region $[-\rho + \rho_0, \rho - \rho_0] \times [0, 1]$ to maps u_1 and u_2, respectively;
- on the growing middle region (4.4), the maps w_j converge to a constant (since there is at most one bubble).

We claim that w_j eventually lies in the neighborhood of $\mathcal{N} \subset N\mathcal{N}$ where the Newton iteration is defined, in other words that the convergence above also holds with respect to the weighted Sobolev norms on preglued domains. An action argument shows that if $\rho > \rho_0 \gg 0$ is large enough, the entire strip $[-\rho+\rho_0, \rho-\rho_0] \times [0,1]$ maps into the coordinate chart around the double point a where $f(\Sigma)$ looks like $\mathbb{R}^n \cup i\mathbb{R}^n$. The resulting holomorphic map to $(\mathbb{C}^n, \mathbb{R}^n \cup i\mathbb{R}^n)$ has a Fourier expansion

$$z \mapsto \sum_{n\in\mathbb{Z}} c_n e^{(n+\frac{1}{2})\pi z}.$$

As in the proof of [5, Theorem 1.3], the C^0-norm near the ends of the strip controls the norm in $\mathcal{E}(w_\rho) \oplus V(w_\rho)$, where $w_\rho = u_1 \#_\rho u_2$. □

5. Constructing a parallelizable bounding manifold

In this section we show that if the stable Gauss map Gf of $f : \Sigma \to \mathbb{C}^n$ is homotopic to that of the Whitney immersion, then $\Sigma \in bP_{n+1}$, i.e. Σ bounds a parallelizable $(n+1)$-manifold. This excludes numerous diffeomorphism types by results of [20]. We obtain the bounding manifold for Σ by attaching a cap to the outer boundary $\mathcal{F}_0^{\mathrm{bd}} = \partial\mathcal{F}_0 - \Sigma \approx \mathcal{F}_1 \times \mathcal{M}$. We do this in the simplest possible way: let

$$\mathcal{B} = \mathcal{F}_0 \cup_{\mathcal{F}_0^{\mathrm{bd}}} (\mathcal{D} \times \mathcal{M})$$

where $\mathcal{D} = \sqcup D^2$ is a finite collection of closed disks abstractly bounding the 1-manifold $\mathcal{F}_1$.

Proposition 5.1. *The space $\mathcal{B}$ admits the structure of a compact C^1-smooth manifold, with boundary canonically diffeomorphic to the homotopy sphere Σ.*

Proof. This is an immediate consequence of Lemma 4.1. □

We claim that, under the homotopy assumption on the Gauss map, the manifold $\mathcal{B}$ constructed above has trivial tangent bundle. Since for manifolds with boundary triviality and stable triviality of the tangent bundle are equivalent, it suffices to show that $T\mathcal{B}$ is trivial in the reduced KO-theory $\widetilde{KO}(\mathcal{B})$.

5.1. **The Whitney immersion.**

To analyze $T\mathcal{B}$, following [11], we use index theory, and aim to trivialize the appropriate index bundles over configuration spaces of smooth maps of punctured disks. For these to be sufficiently explicit as to be amenable to study, we reduce to a model situation.

Definition 5.2. Let $W\colon S^n \to \mathbb{C}^n$ denote the Whitney immersion,

$$W\colon \{(x,y) \in \mathbb{R}^n \times \mathbb{R} \,|\, |x|^2 + y^2 = 1\} = S^n \to \mathbb{C}^n \quad W(x,y) = (1+iy)x.$$

This is a Lagrangian immersion of a sphere with a unique double point, for which the corresponding Reeb chord has Maslov index n. Consider now a homotopy n-sphere Σ and a Lagrangian immersion $f\colon \Sigma \to \mathbb{C}^n$ with one double point a of index n. Fix once and for all a stable framing of the tangent bundle of Σ, which exists by [20], and fix a Hamiltonian isotopy of $\mathbb{C}^n$ that makes f agree with the Whitney immersion in a neighborhood of the double point a.

In this situation, any choice of stable framing η of the tangent bundle TS^n of the standard sphere induces a *difference element* $\delta(\eta) \in U$, obtained as follows. Since f and W have the same Maslov index, the maps may be homotoped (relative to a neighborhood of the double point) so as to co-incide in a neighborhood of a path connecting the double point preimages. The complement of such a path is an n-disk in either S^n or Σ. Since the tangent bundles are framed, the Gauss maps df and dW are naturally valued in $U(n)$, and co-incide on the boundaries of these disks. They therefore patch to define a map $\delta : S^n \to U(n)$, well-defined up to homotopy. The difference element $\delta(\eta)$ is the corresponding stable map $S^n \to U$.

Lemma 5.3. *There exists a pl-isotopy $\Phi_t \colon \Sigma \to \mathbb{C}^n$ fixed around $f(a)$ such that $\Phi_0 = \mathrm{id}$ and $\Phi_1 \circ f$ is a parameterization of the Whitney immersion. Furthermore, if the stable Gauss map Gf is homotopic to the stable Gauss map GW of the Whitney immersion, there is a stable framing η of TS^n so that the induced difference element $\delta(\eta) = 0$.*

Proof. The first statement follows from a PL-isotopy extension theorem due to Hudson and Zeeman [19], compare to [11, Lemma 3.13]. For the second statement, note that changing the framing η on the standard sphere corresponds to precomposing the map δ with a map into O_n. This is sufficient to kill the stable obstruction $\delta(\eta)$ precisely when the stable Gauss maps into U/O are homotopic. □

Corollary 5.4. *If the stable Gauss maps of f and W are homotopic, there is a one-parameter family of C^0-homeomorphisms $\phi_t \colon \mathbb{C}^n \to \mathbb{C}^n$ and a 1-parameter family of continuous maps $\psi_t : S^n \to U$ with the properties that*

1. *ϕ_0 is the identity;*
2. *$\phi_1(f(\Sigma)) = W(S^n)$;*
3. *$\psi_0 = df$ and $\psi_1 = dW \circ \phi_1$.*

5.2. **Index bundles and configuration spaces.** Recall the configuration spaces $\mathcal{X}_{p,q}$ for (Floer) holomorphic disks with boundary on Σ, with p positive and q negative boundary punctures, modeled on Sobolev spaces with two derivatives in L^2. It is standard that the KO-theory class of the tangent bundle of the solution spaces $\mathcal{F}_{p,q}$ and $\mathcal{M}_{p,q}$ of the perturbed and unperturbed $\bar\partial$-equation, respectively, is given by the linearization of the $\overline{\partial}$-operator, which in turn is globally restricted from the ambient configuration space

$$\left[T\mathcal{X}_{p,q} \xrightarrow{D(\bar\partial)} \mathcal{Y}_{p,q} \right],$$

where $\mathcal{Y}_{p,q}$ is the Sobolev space of complex anti-linear maps. More precisely, for a Lagrangian immersion $L \to \mathbb{C}^n$, any map $D_{p,q} \to \mathbb{C}^n$ with boundary on L defines a $\overline{\partial}$-operator in a trivial bundle over the disk. This is a formally self-adjoint Fredholm operator, and hence there is a polarized Hilbert bundle (polarized by the positive / negative spectrum) over the configuration space of all smooth disk maps. Any such bundle is classified by a map

$$\mathrm{Maps}((D_{p,q}, \partial D_{p,q}), (\mathbb{C}^n, L)) \longrightarrow BGL_{res} \simeq U/O \tag{5.1}$$

into the restricted Grassmannian of Hilbert space, which by [26] is homotopy equivalent to U/O. (The induced map on constants is exactly the stable Gauss map of the immersion.) (5.1) tautologically classifies the index bundle of the $\overline{\partial}$-operator over configuration space. By conjugating with the formal isotopy of Lagrangian planes produced by the final part of Corollary

5.4, the maps (5.1) associated to the immersion f of Σ and the Whitney immersion are homotopic. This proves:

Corollary 5.5. *The index bundle on the configuration space $\mathcal{X}_{p,q}$ is stably trivial if and only if the index bundle on the corresponding configuration space for the Whitney immersion is stably trivial.* □

We restrict attention to the configuration spaces relevant to our study and write $\mathcal{X}_0 = \mathcal{X}_{0,0}$, $\mathcal{X}_1 = \mathcal{X}_{0,1}$, and $\mathcal{X} = \mathcal{X}_{1,0}$ for the configuration spaces that contain $\mathcal{F}_0$, $\mathcal{F}_1$, and $\mathcal{M}$, respectively.

For a compact connected cell complex K and $x, y \in K$ let $\Omega(K; x, y)$ denote the space of paths in K from x to y, which is homotopy equivalent to the based loop space $\Omega(K) = \Omega(K; x, x)$ of K. Let $\mathcal{L}(K)$ denote the free loop space of K. Recall that $\{a^+, a^-\} \subset \Sigma$ are the preimages of the double point $a \in \mathbb{C}^n$ of the immersion f.

Lemma 5.6. *There are homotopy equivalences*

$$\mathcal{X}_1 \simeq S^1 \times \Omega(\Sigma; a^-, a^+); \quad \mathcal{X} \simeq \Omega(\Sigma; a^+, a^-); \quad \mathcal{X}_0 \simeq \mathcal{L}(\Sigma),$$

In particular, the homotopy types of the configuration spaces are independent of the smooth structure on the homotopy sphere Σ.

Proof. In all cases, the homotopy fiber of the restriction to the boundary is contractible via linear homotopies. Note that the S^1-factor encodes the location of the boundary puncture of maps in $\mathcal{X}_1$. The final statement follows since the homotopy type of the based and free loop spaces is a homotopy invariant of compact cell complexes. □

We write I_0, I_1, and I to denote the index bundles over $\mathcal{X}_0$, $\mathcal{X}_1$, and $\mathcal{X}$ respectively. Let $Q \to U\Sigma$ denote the vertical tangent bundle to the unit sphere bundle $U\Sigma \subset T\Sigma$, and $\widehat{Q}$ the fiberwise Thom space of Q, so the fiber $\widehat{Q}_\sigma$ of $\widehat{Q}$ over $\sigma \in \Sigma$ is the Thom space of the tangent bundle $T(U_\sigma\Sigma)$ of the fiber of the unit sphere bundle at σ. Note that the Thom space contains a distinguished point ∞. Let

$$\Sigma \vee \widehat{Q}_\sigma = \Sigma \cup \widehat{Q}_\sigma / \sigma \sim \infty.$$

Lemma 5.7. *There are embeddings*

$$\iota\colon S^{n-1} \to \Omega(\Sigma; a^\pm, a^\mp) \qquad \text{and} \qquad \kappa\colon \Sigma \vee \widehat{Q}_\sigma \ \to \mathcal{L}(\Sigma)$$

which give a $2n-3$ respectively $3n-5$ skeleton for the given spaces. In other words ι is $(2n-3)$-connected and κ is $(3n-5)$-connected.

Proof. The skeleta arise from the usual Morse-Bott theory of the energy functional for the round metric on the standard sphere S^n. For the path space $\Omega(\Sigma; a^\pm, a^\mp)$, take a^+ and a^- to be antipodal points. Then the energy functional is Bott-degenerate with Bott-manifolds diffeomorphic to S^{n-1} of indices $2k(n-1)$, $k = 0, 1, 2, \ldots$. It follows that the Bott-manifold of index 0 gives a $(2n-3)$-skeleton.

For the free loop space, consider the fibration $\mathcal{L}(\Sigma) \to \Sigma$ with fiber the based loop space $\Omega(\Sigma)$. Again the energy functional is Bott-degenerate with critical manifold Σ of index 0 corresponding to constant loops, and critical manifolds $U\Sigma$ of indices $2k(n-1)$, $k = 1, 2, 3, \ldots$. The closure of the unstable manifold of the index $2(n-1)$ critical manifold $U\Sigma$ gives an embedding of $\widehat{Q}$, see [11, Lemma 7.1], which then gives a $(3n-3)$-skeleton. Removing the fiber over one point we get a $(3n-4)$-skeleton homotopy equivalent to $\Sigma \vee \widehat{Q}_\sigma$. □

Combining Lemmas 5.6 and 5.7 we get a corresponding $(2n-3)$-skeleton $S^{n-1} \approx K^{2n-3} \subset \mathcal{X}$, a $(2n-2)$-skeleton $S^1 \times S^{n-1} \approx K_1^{2n-2} \subset \mathcal{X}_1$, and a $(3n-4)$-skeleton $\Sigma \vee \widehat{Q}_\sigma \approx K_0^{3n-4} \subset \mathcal{X}_0$, consisting of disk maps with boundary paths in $\iota(S^{n-1})$, with boundary paths in $\iota(S^{n-1})$ and base point according to the S^1-factor, and with boundary loops in $\kappa(\widehat{Q})$, respectively.

Lemma 5.8. *The index bundles I_0, I, and I_1 are stably trivial over the skeleta K_0^{3n-4}, K^{2n-3}, and K_1^{2n-2}, respectively.*

Proof. By Corollary 5.5, it is sufficient to find stable trivializations of the index bundles for the Whitney immersion, and we can moreover assume that we work with the undeformed Cauchy-Riemann equation and the standard $\bar{\partial}$-operator. We use a model of the Whitney sphere that is almost flat, i.e. for some small $\epsilon > 0$, we scale $\mathbb{R}^n$ and $i\mathbb{R}^n$ in the model of Definition 5.2 by ϵ^{-1} and ϵ, respectively.

Consider first $I \to K^{2n-3}$. The Lagrangian boundary condition corresponding to a geodesic connecting a^+ to a^- splits into a 1-dimensional problem, corresponding to the tangent line of the curve in Figure 1, of index 2 with 2-dimensional kernel, and an $(n-1)$-dimensional problem with almost constant boundary conditions and hence with $(n-1)$-dimensional kernel. It follows that the index bundle is stably equivalent to TS^{n-1}, in particular stably trivial as claimed.



Figure 1. The Lagrangian boundary condition along a disk with a positive puncture. *Upper*: the front projection of the Whitney sphere, representing the immersion as a multi-sheeted graph (one recovers $y_j = dz/dx_j$). *Lower*: the intersection of the Whitney sphere and a complex co-ordinate line in $\mathbb{C}^n$.

For $I_1 \to K_1^{2n-2}$ the boundary condition splits in the same way, except now the 1-dimensional problem corresponds to the curve in Figure 1 oriented in the opposite direction, since the puncture is negative. The index for that problem now equals -1, whilst the $\bar{\partial}$-operator on the almost constant $(n-1)$-dimensional orthogonal boundary condition is an isomorphism. Thus, the index bundle is trivial.

Finally, consider $I_0 \to K_0^{3n-4}$. Restricted to constant loops in S^n, the index bundle is isomorphic to the tangent bundle, hence is stably trivial. Consider now $\widehat{Q}_\sigma$. Taking $\sigma = a^+$ we find that the geodesic loops are unions of the paths considered above. Using sufficiently stretched disks of the form $H_1 \#_\rho H_2$, the boundary condition over $U_\sigma S^n$ can be considered as a sum of the two problems studied above. Stabilizing the operator defining I_1 once, the corresponding operator becomes an isomorphism. By considering an explicit model of the Morse flow, compare to the proof of [11, Lemma 7.3], one sees that the restriction of I_0 to $\widehat{Q}_\sigma = MT(U_\sigma S^n)$ is stably isomorphic to the pull-back of the tangent bundle $T(U_\sigma S^n)$, hence is stably trivial. The trivialization can clearly be glued to that over S^n at σ, which completes the proof. □

5.3. **Coherent stable trivializations.**

Consider stable trivializations Z_0, Z_1, and Z of I_0, I_1, and I, respectively. Recall the pre-gluing map from Section 4.1. For any compact subsets $B_1 \subset \mathcal{X}_1$ and $B \subset \mathcal{X}$ there is an analogous map

$$\mathrm{PG}_\rho \colon B_1 \times B \to \mathcal{X}_0$$

for all sufficiently large ρ. Suppose now we stabilize the linearized $\bar{\partial}$-operators defining I_1 and I over B_1 and B, by adding trivial bundles $\mathbb{R}^{m_1}$ and $\mathbb{R}^m$ to make the operators everywhere surjective. For ρ large enough, there is a linear gluing isomorphism

$$G_\rho \colon \ker(D\bar{\partial} \oplus \mathbb{R}^{m_1})_{u_1} \oplus \ker(D\bar{\partial} \oplus \mathbb{R}^m)_{u_2} \to \ker(D\bar{\partial} \oplus \mathbb{R}^{m_1} \oplus \mathbb{R}^m)_{u_1 \#_\rho u_2} \tag{5.2}$$

that takes kernel elements to cut-off versions supported in the respective halves of the preglued domain, see [11, Lemma 7.2].

We say that stable trivializations (Z_1, Z, Z_0) are (d_1, d)-coherent if for any compact CW-complexes $B_1 \subset \mathcal{X}_1$ and $B \subset \mathcal{X}$ of dimensions d_1 and d, respectively, the stable trivializations $\mathrm{PG}_\rho^* Z_0$ and $G_\rho(Z_1 \oplus Z)$ of the pull-back of the index bundle $\mathrm{PG}^* I_0$ are homotopic. Coherent trivializations generalize the more familiar notion of coherent orientations.

Lemma 5.9. *For any stable trivialization Z_0 of I_0 and Z_1 of I_1, with Z_1 trivial over the 1-skeleton (i.e. over the S^1-factor in $K_1^{2n-2} \approx S^1 \times S^{n-1}$), there is a stable trivialization Z of I such that (Z_1, Z, Z_0) is $(1, n-1)$ coherent.*

Proof. After homotopy, we may assume that any closed 1-complex B_1 mapping to $\mathcal{X}_1$ maps into the 1-skeleton $S^1 \times v$, for some $v \in S^{n-1}$, and that any compact $(n-1)$-complex mapping to $\mathcal{X}$ maps into $K^{2n-1} \approx S^{n-1}$. Recall from Lemma 5.8 that the trivialization Z_0 of I_0 over K_0^{3n-4} is constructed using almost broken disks. By definition, in such a trivialization $Z_0 = G(Z_1 \oplus Z)$, where we stabilize the operator defining I_1 once to make it an isomorphism over S^{n-1}, and where Z is some given trivialization of I. □

5.4. **Proof of Theorem 1.1.**

Lemma 2.2 shows that K is a homotopy sphere. The regular homotopy assumption together with Lemma 5.8 implies that the index bundles I_1, I_0, and I are stably trivial, and using Lemma 5.9 we find a coherent triple (Z_1, Z, Z_0) of stable trivializations where Z_1 is trivial over the 1-skeleton. Note that the tangent bundle of the 1-manifold $\mathcal{F}_1$ is canonically isomorphic to I_1. Hence, our assumption on the stable trivialization Z_1 over the 1-skeleton implies that the induced stable trivialization of $T\mathcal{F}_1$ extends to a trivialization $\tilde{Z}_1$ of $T\mathcal{D}$. We thus get a stable trivialization $\tilde{Z}_1 \oplus Z$ of $T(\mathcal{D}\times\mathcal{M})$, which near $\partial^{\mathrm{bd}}\mathcal{F}_0$ agrees with $G(Z_1 \oplus Z)$ and hence extends as Z to a stable trivialization of $T\mathcal{F}_0$. We conclude that $T\mathcal{B}$ is stably trivial. □

5.5. **Signature.** Any (not necessarily closed) $4k$-manifold X has a signature $\sigma(X)$ which is the number of positive minus the number of negative eigenvalues of the intersection form on $H_{2k}(X;\mathbb{Q})$. According to [20], the exotic nature of a $(4k-1)$-dimensional homotopy sphere is reflected in the possible signatures of parallelizable manifolds which it bounds.

Now suppose, as in previous sections, that $H\colon \mathbb{C}^n \times [0,1] \to \mathbb{R}$ is a compactly supported Hamiltonian function whose time 1 flow displaces a Lagrangian immersion $f\colon \Sigma \to \mathbb{C}^n$ with one double point of index n. Let $\mathcal{B}_H$ denote the bounding manifold, with $\partial\mathcal{B}_H = \Sigma$, obtained from capping the space of Floer solutions as usual, so $\mathcal{B}_H = \mathcal{F}_0 \cup_{\mathcal{F}_1\times\mathcal{M}} (\mathcal{D}\times\mathcal{M})$.

Proposition 5.10. *$\sigma(\mathcal{B}_H)$ is independent of H, hence defines an invariant $\sigma(f)$ of the original Lagrangian immersion $f:\Sigma\to\mathbb{C}^n$. If $W : S^n \to \mathbb{C}^n$ is the Whitney immersion, then $\sigma(W) = 0$.*

Sketch. Both statements are applications of Novikov additivity; we just sketch the main ideas. The first result is obtained by considering a family of displacing Hamiltonians $\{H_s\}$ interpolating two given such H_0, H_1, and a corresponding family of Floer-equations $(3.3)_R$ parametrized by $R \in \mathfrak{h}$, a half-plane, see Figure 2. The Gromov-Floer compactified space of Floer disks with no punctures over $\mathfrak{h}$ admits a cap of the shape $S\times\mathcal{M}$ for a solid handlebody S, and shows that $\mathcal{B}_{H_0} \cup_\Sigma \mathcal{B}_{H_1}$ is a global boundary, hence has trivial signature. (For this argument, C^1-structures and parallelizability play no role.) The second result follows from a similar construction, using the fact that the Whitney immersion of the n-sphere extends to an exact Lagrangian immersion of an $(n+1)$-ball. We leave details to the reader. □



Figure 2. Cauchy-Riemann problems parametrized by a half-plane

There is a natural evaluation map $ev_1 : \mathcal{F}_0 \to \mathbb{C}^n$ which extends a given immersion f, but in general there is no reason to expect F to be better-behaved than a generic map. The regular homotopy classes of Lagrangian immersions in Euclidean space of a homotopy $(8k-1)$-sphere are classified by $\pi_{8k-1}(U) \cong \mathbb{Z}$. The set of elements realized by immersions with one double point of some given homotopy sphere Σ may be related, via the signature of the bounding manifold $\mathcal{B}$, to the singularities of the evaluation map ev_1 on high-dimensional moduli spaces of Floer disks. It would be interesting to understand if those singularities can be interpreted in terms of the local analysis of the underlying Floer disks themselves.

References

[1] M. Abouzaid. *Framed bordism and Lagrangian embeddings of exotic spheres.* Ann. of Math. **175** (2012) 71–185.

[2] M. Abouzaid. *Nearby Lagrangians with vanishing Maslov class are homotopy equivalent.* Invent. Math. **189** (2012) 251–313. 2.1

[3] F. Bourgeois, T. Ekholm and Y. Eliashberg. *Effect of Legendrian surgery.* Geom. Topol. **16** (2012) 301–389. 2

[4] M. Damian. *Floer homology on the universal cover, a proof of Audin's conjecture, and other constraints on Lagrangian submanifolds.* Comment. Math. Helv. **87** (2012) 433–462. 1, 2, 2

[5] T. Ekholm. *Morse flow trees and Legendrian contact homology in 1-jet spaces.* Geom. Topol. **11** (2007) 1083–1224. 4.4

[6] T. Ekholm, J. Etnyre and J. Sabloff, *A duality exact sequence for Legendrian contact homology*, Duke Math. J. **150** (2009) 1–75. 2, 2, 2

[7] T. Ekholm, J. Etnyre and M. Sullivan, *Orientations in Legendrian contact homology and exact Lagrangian immersions*, Internat. J. Math. **16** (2005) 453–532.

[8] T. Ekholm, J. Etnyre and M. Sullivan. *The contact homology of Legendrian submanifolds in* $\mathbb{R}^{2n+1}$. J. Differential Geom. **71** (2005) 177–305. 2, 3.2

[9] T. Ekholm, J. Etnyre and M. Sullivan. *Non-isotopic Legendrian submanifolds in* $\mathbb{R}^{2n+1}$. J. Differential Geom. **71** (2005) 85–128.

[10] T. Ekholm, J. Etnyre and M. Sullivan. *Legendrian contact homology in* $P \times \mathbb{R}$. Trans. Amer. Math. Soc. **359** (2007) 3301–3335. 2, 2, 3.2, 3.3, 4.3

[11] T. Ekholm and I. Smith, *Exact Lagrangian immersions with a single double point*, Preprint, arXiv:1111.5932 (document), 1, 1, 3.1, 4, 4, 4.1, 4.4, 4.4, 5.1, 5.1, 5.2, 5.2, 5.3

[12] T. Ekholm, Y. Eliashberg, E. Murphy and I. Smith. *Manuscript in preparation.* 1

[13] Y. Eliashberg and E. Murphy. *Manuscript in preparation.* 1

[14] A. Eriksson-Östman. *Manuscript in preparation.* 1, 2

[15] A. Floer. *Morse theory for Lagrangian intersections.* J. Differential Geom. **28** (1988) 513–547.

[16] A. Floer. *Monopoles on asymptotically flat manifolds.* The Floer memorial volume, Progr. Math., **133** (1995) 3–41, Birkhäuser, Basel. 4

[17] M. Gromov. *Pseudoholomorphic curves in symplectic manifolds.* Invent. Math. **82** (1985) 307–347. 1

[18] M. Gromov. *Partial differential relations.* (1986) Springer.

[19] J. F. Hudson and E. C. Zeeman. *On combinatorial isotopy.* Inst. Hautes Etudes Sci. Publ. Math. **19** (1964) 69–94. 5.1

[20] M. Kervaire and J. Milnor. *Groups of homotopy spheres, I.* Ann. of Math. **77** (1967) 504–537. 1, 5, 5.1, 5.5

[21] D. Kwon and Y.-G. Oh. *Structure of the image of pseudo-holomorphic discs with totally real boundary condition.* Comm. Anal. Geom. **1** (2000) 31–82.

[22] L. Lazzarini. *Existence of a somewhere injective pseudo-holomorphic disc.* Geom. Funct. Anal. **10** (2000) 829–862.

[23] E. Murphy. *Loose Legendrian embeddings in high-dimensional contact manifolds.* Preprint, available at arXiv:1201.2245. 1

[24] Y.-G. Oh. *Gromov-Floer theory and disjunction energy of compact Lagrangian submanifolds.* Math. Res. Lett. **6** (1997) 895–905. 3.1

[25] Y.-G. Oh. *Floer cohomology of Lagrangian intersections and pseudo-holomorphic disks, I.* Comm. Pure Appl. Math. **46** (1993) 949–993. 3.3

[26] A. Pressley and G. Segal. *Loop groups.* (1986) Oxford University Press. 5.2

[27] D. Sauvaget, *Curiosités Lagrangiennes en dimension 4* Ann. Inst. Fourier (Grenoble) **54** (2004) 1997–2020. 1

Department of mathematics Uppsala University, Box 480, 751 06 Uppsala, Sweden
Institut Mittag-Leffler, Aurav. 17, 182 60 Djursholm, Sweden

E-mail address: `tobias.ekholm@math.uu.se`

Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, CB3 0WB, UK

E-mail address: `is200@cam.ac.uk`